OMIM







Taxonomy

Clipboard

PubMed

Nucleotide

Q PubMed Search

Genome Structure Protein

Clear Go

Details

8

About Entrez

Display

for

Limits

Sort Abstract

Preview/Index



PopSet

History

Order

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Browser Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Gateway TOXNET** Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Privacy Policy

□ 1: Cancer Lett 2002 Feb 8;176(1):47-55

Related Articles, Links

ELSEVIER SCIENCE FULL-TEXT ARTICLE

Ectopic expression of homeodomain protein CDX2 in intestinal metaplasia and carcinomas of the stomach.

Bai YQ, Yamamoto H, Akiyama Y, Tanaka H, Takizawa T, Koike M, Kenji Yagi O, Saitoh K, Takeshita K, Iwai T, Yuasa Y.

Department of Surgery, Graduate School of Medicine and Dentistry, Tokyo Medical and Dental University, Tokyo, Japan.

The roles of CDX2 and CDX1 homeobox genes during gastric carcinogenesis remain poorly defined. We have studied the expression of CDX2/1 in gastric cancers and intestinal metaplasia (IM) of 69 gastric carcinoma patients by immunohistochemistry. CDX2/1 were shown to be ectopically overexpressed in IM in 41 (85%) of 48, and 47 (90%) of 52 cases, respectively. The expression of CDX2/1 was detected in 38 (55%) and 51 (74%) of the 69 gastric carcinomas, respectively. The histological type of the gastric carcinomas was independently associated with CDX2 expression, but not with that of CDX1, with higher CDX2 expression in intestinal type (differentiated type) than in diffuse type (undifferentiated type) gastric carcinomas. Our results thus suggest that CDX2 and CDX1 may play a role during IM formation and gastric carcinogenesis.

PMID: 11790453 [PubMed - indexed for MEDLINE]

Text Abstract Sort Display

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services